

VARIANCE OF LIPSCHITZ FUNCTIONS AND AN ISOPERIMETRIC PROBLEM FOR A CLASS OF PRODUCT MEASURES*

S.G. Bobkov[†] and C. Houdré[‡]

July 10, 1995

Abstract

The maximal variance of Lipschitz functions (with respect to the ℓ_1 -distance) of independent random vectors is found. This is then used to solve the isoperimetric problem, uniformly in the class of product probability measures with given variance.

1 Statements

Let $\xi = (\xi_1, \dots, \xi_n)$ be a vector of independent random variables with finite variance $\sigma_i^2 = \mathbf{Var} \xi_i$, $1 \leq i \leq n$. Denote by \mathcal{F}_1 the class of all functions on \mathbf{R}^n which are Lipschitz with respect to the ℓ^1 -distance

$$d_1(x, y) = \|x - y\|_1 = \sum_{k=1}^n |x_k - y_k|, \quad x, y \in \mathbf{R}^n.$$

By definition, $f \in \mathcal{F}_1$, if for all $x, y \in \mathbf{R}^n$, $|f(x) - f(y)| \leq d_1(x, y)$. Let $S_n = \xi_1 + \dots + \xi_n$.

*Key Words: Isoperimetry, Lipschitz function, Variance inequality

[†]Research supported in part by the ISF grant NZX000 and NZX300

[‡]Research supported in part by an NSF Mathematical Sciences Postdoctoral Fellowship

Theorem 1 *In the class \mathcal{F}_1 , the maximal value of $\mathbf{Var}f(\xi)$ is attained at the function $f(x) = x_1 + \dots + x_n$. In other words, for any $f \in \mathcal{F}_1$,*

$$\mathbf{Var}f(\xi) \leq \mathbf{Var}S_n = \sum_{i=1}^n \sigma_i^2. \quad (1.1)$$

X.Fernique [F, Théorème 3.2] proved an inequality similar to (1.1) for $f \in \mathcal{F}_1$, convex. However, ξ there, is only assumed to be symmetrically distributed, i.e., for all $\varepsilon_i = \pm 1$, the random vectors $(\varepsilon_1 \xi_1, \dots, \varepsilon_n \xi_n)$ have the same distribution (of course, this is fulfilled if the ξ_i are i.i.d. with a symmetric one-dimensional distribution). In contrast to the difficult proof of Fernique, Theorem 1 can easily be obtained by induction.

Theorem 1 also has the following consequence: Denote by $M^n(\sigma)$ the family of all the product measures $\mu = \mu_1 \otimes \dots \otimes \mu_n$ on \mathbf{R}^n with given variance $\mathbf{Var}(\mu) = \sigma^2$, where

$$\mathbf{Var}(\mu) = \sum_{i=1}^n \int_{\mathbf{R}} \left| x - \int_{\mathbf{R}} t d\mu_i(t) \right|^2 d\mu_i(x).$$

Hence, with the above notations, $\mathbf{Var}(\mu) = \mathbf{Var}S_n$. Now, given a set $A \subset \mathbf{R}^n$ and $h > 0$, denote by

$$A^h = A + hB_1 = \{x \in \mathbf{R}^n : d_1(a, x) < h, \text{ for some } a \in A\}$$

the open h -neighbourhood of A (B_1 is the open ℓ^1 -unit ball in \mathbf{R}^n). From Theorem 1 we obtain a solution to the isoperimetric problem with respect to the ℓ^1 -distance uniformly in the class $M^n(\sigma)$ controlled by the parameter σ .

Theorem 2 *For any $h > 0$, $\sigma > 0$ and $p \in (0, 1)$,*

$$\inf_{\mu \in M^n(\sigma)} \inf_{\mu(A) \geq p} \mu(A^h) = \begin{cases} p, & \text{if } h \leq \frac{\sigma}{\sqrt{p(1-p)}}, \\ 1 - \frac{p\sigma^2}{ph^2 - \sigma^2}, & \text{if } h \geq \frac{\sigma}{\sqrt{p(1-p)}}. \end{cases} \quad (1.2)$$

The first infimum in (1.2) is taken over all the $\mu \in M^n(\sigma)$, the second is taken over all the Borel sets A of μ -measure greater or equal to p . In particular, from Theorem 2, we have:

Corollary 3 *Given $\sigma > 0$ and $p \in (0, 1)$, one can guarantee that $\mu(A^h) > p$ regardless of the dimension $n \geq 1$, regardless of the measure $\mu \in M^n(\sigma)$, and regardless of the set $A \subset \mathbf{R}^n$ of μ -measure greater or equal to p , if and only if*

$$h > h(p, \sigma) \equiv \frac{\sigma}{\sqrt{p(1-p)}}.$$

Otherwise, it is possible to have $\mu(A^h) = p$.

Equality in (1.2) is already attained when $n = 1$. Indeed, denote by δ_x the unit mass at the point $x \in \mathbf{R}$. If $h \leq h(p, \sigma)$, take

$$\mu = p\delta_0 + (1-p)\delta_{h(p,\sigma)}, \quad A = \{0\}.$$

Then, $\mathbf{Var}(\mu) = \sigma^2$, $A^h = (-h, h)$, so $\mu(A^h) = p = \mu(A)$. If $h \geq h(p, \sigma)$, take

$$\mu = p\delta_0 + q\delta_x + r\delta_h, \quad A = \{0\},$$

with $r = p\sigma^2/(ph^2 - \sigma^2)$, $q = 1 - p - r$, $x = rh/(p + q)$. Then, it is again easy to verify that $\mathbf{Var}(\mu) = \sigma^2$, and that $\mu(A^h) = 1 - p\sigma^2/(ph^2 - \sigma^2)$.

Since equality in (1.2) is attained when $n = 1$, (1.2) will not change if the h -neighbourhood is defined with respect to the ℓ^2 -distance, or more generally, with respect to the ℓ^α -distance in \mathbf{R}^n , $1 \leq \alpha \leq +\infty$. Indeed, the ℓ^α -unit ball B_α is larger than B_1 , hence, $A + hB_1 \subset A + hB_\alpha$, and therefore $\mu(A + hB_1) \leq \mu(A + hB_\alpha)$. Hence, the same inequality holds when one takes the second infimum in (1.2). But, all the balls B_α coincide when $n = 1$ (in which case equality in (1.2) is attained).

For individual measures μ (for example, for those having finite exponential moments) there exist estimates for $1 - \mu(A^h)$ which decrease exponentially when $h \rightarrow +\infty$ (see M.Talagrand [T2]). For example, given $\sigma_i > 0$, $1 \leq i \leq n$, let $\mu = \mu_1 \otimes \cdots \otimes \mu_n \in M^n(\sigma)$, where $\mu_i = (\delta_{\sigma_i} + \delta_{-\sigma_i})/2$, $\sigma^2 = \sigma_1^2 + \cdots + \sigma_n^2$. Then, as shown in [T2, Proposition 2.1.1, Theorem 2.4.1] (see also M.Ledoux [L, p.24] for an extension to non-identical marginals), if $h > 0$, $\mu(A) = p$, then

$$\mu(A + hB_1) \geq 1 - \frac{1}{p} \exp(-h^2/4\sigma^2). \quad (1.3)$$

When all the $\sigma_i = 1$, the extremal sets minimizing $\mu(A^h)$, while $\mu(A) = p$ is fixed, are known and were obtained by L.H.Harper [H]. If one minimizes

$\mu(A^h)$ over all convex sets A , the situation changes considerably, and we then deal with a much more powerful concentration principle discovered by M. Talagrand (see [T1],[T2]). In particular, when all the $\sigma_i = 1$, one has

$$\mu(A + hB_2) \geq 1 - \frac{1}{p} \exp(-h^2/8).$$

In our case, since one looks for a uniformly minimal value of $\mu(A + hB_1)$, it does not matter whether one considers convex sets or arbitrary sets, since the extremal $A = \{0\}$ is convex.

To finish this section, we give an inequality which is actually equivalent to the second part of (1.2). For non-empty sets $A, B \subset \mathbf{R}^n$, let $d_1(A, B) = \inf\{d_1(a, b) : a \in A, b \in B\}$.

Corollary 4 *For any $\mu \in M^n(\sigma)$, and any non-empty Borel sets $A, B \subset \mathbf{R}^n$,*

$$d_1(A, B) \leq \sigma \sqrt{\frac{1}{\mu(A)} + \frac{1}{\mu(B)}}. \quad (1.4)$$

Let $\mu(A) > 0$, $\mu(B) > 0$ be such that $\mu(A) + \mu(B) \leq 1$. Then, choosing $B = \{h\}$ with h equal to the right-hand side of (1.4), it is easily seen that equality in (1.4) is attained at the same measure μ and the same set $A = \{0\}$ as the second inequality in (1.2).

2 Proofs

A statement slightly more general than Theorem 1 will actually be proved. Assume we have n measurable spaces (X_k, Σ_k) and n measurable functions $h_k = h_k(x_k, y_k)$ defined on $X_k \times X_k$, $1 \leq k \leq n$, and which vanish on the diagonal $x_k = y_k$. Let ξ_k be independent random variables with values in X_k , $1 \leq k \leq n$, such that

$$2\sigma_k^2 = \mathbf{E} h_k^2(\xi_k, \eta_k) < +\infty,$$

where η_k is an independent copy of ξ_k . Put $\xi = (\xi_1, \dots, \xi_n)$.

Lemma 5 *Let f be a measurable function defined on $X_1 \times \dots \times X_n$ such that*

$$|f(x) - f(y)| \leq \sum_{k=1}^n h_k(x_k, y_k), \quad (2.1)$$

for $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in X_1 \times \dots \times X_n$. Then,

$$\mathbf{Var} f(\xi) \leq \sum_{k=1}^n \sigma_k^2. \quad (2.2)$$

Proof. This lemma is proved by induction on the dimension n . For $n = 1$, and since $2\mathbf{Var} f(\xi) = \int \int (f(\xi) - f(\eta))^2 d\mu(\xi) d\mu(\eta)$, (2.2) is immediate. Assume now that (2.2) is true for n . Denote by μ_{n+1} the distribution of ξ_{n+1} , and by P_n the distribution of the random vector (ξ_1, \dots, ξ_n) , thus $P_{n+1} = P_n \otimes \mu_{n+1}$ is the distribution of $(\xi_1, \dots, \xi_{n+1})$. Let $f : X_1 \times \dots \times X_{n+1} \rightarrow \mathbf{R}$ satisfy (2.1). Now, fix x_{n+1} . Since the function $g(x_1, \dots, x_n) = f(x_1, \dots, x_n, x_{n+1})$ satisfies (2.1), making use of the induction hypotheses and writing (2.2) for g we get:

$$\int g^2 dP_n \leq \left(\int g dP_n \right)^2 + \sum_{k=1}^n \sigma_k^2. \quad (2.3)$$

The function $m(x_{n+1}) = \int g dP_n$ is well-defined, measurable and as a function of one variable,

$$|m(x_{n+1}) - m(y_{n+1})| \leq h_{n+1}(x_{n+1}, y_{n+1}).$$

Thus m satisfies (2.1), hence

$$\int m^2 d\mu_{n+1} \leq \left(\int m d\mu_{n+1} \right)^2 + \sigma_{n+1}^2. \quad (2.4)$$

Integrating (2.3) over X_{n+1} (with respect to μ_{n+1}), and taking into account (2.4), gives (2.2) for f . Lemma 5 and thus Theorem 1 are proved.

Proof of Theorem 2

Let $A \subset \mathbf{R}^n$ be such that $\mu(A) \geq p$. Since the function $f(x) = \inf_{a \in A} d_1(a, x)$ belongs to \mathcal{F}_1 , we have by Theorem 1, that $\mathbf{Var} f \leq \sigma^2$. In addition, $f \geq 0$ and $\mu(f = 0) \geq p$. Note also that $A^h = \{x \in \mathbf{R}^n : f(x) < h\}$. To get (1.2), it just remains to appeal to the following result:

Lemma 6 For any $h > 0$, $\sigma > 0$ and $p \in (0, 1)$,

$$\sup \mathbf{P}(\xi \geq h) = \begin{cases} 1 - p, & \text{if } h \leq \frac{\sigma}{\sqrt{p(1-p)}}, \\ \frac{p\sigma^2}{ph^2 - \sigma^2}, & \text{if } h \geq \frac{\sigma}{\sqrt{p(1-p)}}, \end{cases} \quad (2.5)$$

where the supremum is taken over all non-negative random variables ξ on a probability space $(\Omega, \mathcal{B}, \mathbf{P})$ such that $\mathbf{P}(\xi = 0) \geq p$ and $\mathbf{Var} \xi \leq \sigma^2$.

Proof. Denote by $\mathcal{L}(\xi)$ the distribution of ξ . The cases of equality in (2.5) were, in fact, already settled in Section 1. If

$$h \leq h(p, \sigma), \quad \mathcal{L}(\xi) = p\delta_0 + (1-p)\delta_{h(p, \sigma)},$$

then $\mathbf{Var}(\xi) = \sigma^2$, $\mathbf{P}(\xi = 0) = p$, $\mathbf{P}(\xi \geq h) = 1 - p$. If

$$h > h(p, \sigma), \quad \mathcal{L}(\xi) = p\delta_0 + q\delta_x + r\delta_h,$$

where $r = p\sigma^2/(ph^2 - \sigma^2)$, $q = 1 - p - r$, $x = rh/(p+q)$, then as easily verified, we have $\mathbf{Var}(\xi) = \sigma^2$, $\mathbf{P}(\xi = 0) = p$, and $\mathbf{P}(\xi \geq h) = r = p\sigma^2/(ph^2 - \sigma^2)$. So, one needs only to show that whenever $h \geq h(p, \sigma)$,

$$\mathbf{P}(\xi \geq h) \leq \frac{p\sigma^2}{ph^2 - \sigma^2}. \quad (2.6)$$

To prove this last statement, several steps will be needed.

Step 1 (reduction to the bounded case).

It is easy to see, via a usual truncation argument, that ξ can be assumed to be bounded, say, by a constant C . So, $\mathcal{L}(\xi)$ is assumed to be a probability measure F on $[0, C]$.

Step 2 (reduction to discrete case with only four atoms).

Fixing

$$p_0 = \mathbf{P}(\xi = 0) = F(\{0\}), \quad a = \mathbf{E}\xi = \int_0^C x dF(x), \quad b = \mathbf{E}\xi^2 = \int_0^C x^2 dF(x),$$

defines a subset $Z(p_0, a, b)$ of the family Z of all probability measures on $[0, C]$. Then, observe the following:

1) Z is a (convex, compact for the topology of weak convergence) simplex in the space M of all signed measures on $[0, C]$, and the extremal points of Z are just the unit masses δ_x , $0 \leq x \leq C$.

2) $Z(p_0, a, b)$ is the intersection of Z with three hyperspaces $H_0 = \{F \in M : F(\{0\}) = p_0\}$, $H_1 = \{F \in M : \int_0^C x dF(x) = a\}$ and $H_2 = \{F \in M : \int_0^C x^2 dF(x) = b\}$.

3) Therefore, noting that H_1 and H_2 are closed, while $\text{clos}(H_0) = \{F \in M : F(\{0\}) \geq p_0\}$, one easily concludes that the extremal points of $Z(p_0, a, b)$ are just the measures of the form $F = p'_0\delta_0 + p_1\delta_{x_1} + p_2\delta_{x_2} + p_3\delta_{x_3}$, with $p'_0 \geq p_0$.

4) The functional $F \longrightarrow F([h, +\infty))$ is linear and, since the set $[h, +\infty)$ is closed, this functional is upper semi-continuous on Z . Hence, its maximum on $Z(p_0, a, b)$ is attained at an extremal point of $Z(p_0, a, b)$.

Of course, the above maximum should be further maximized over all possible p_0, a, b (in particular, we should have $p_0 \geq p$, $a^2 \leq b$, $b - a^2 \leq \sigma^2$). But in the sequel, one needs only consider the ξ taking four values, and moreover, one may also forget about the condition $\xi \leq C$.

Step 3 (reduction to the four atoms case, one of them being h).

So, by the previous step, it suffices to prove (2.6) when

$$\mathcal{L}(\xi) = p_0\delta_0 + p_1\delta_{x_1} + p_2\delta_{x_2} + p_3\delta_{x_3},$$

$p_0 \geq p$, $p_1, p_2, p_3 \geq 0$, $p_0 + p_1 + p_2 + p_3 = 1$, $x_1, x_2, x_3 \geq 0$. Thus we need to maximize the functional

$$J = p_1 \mathbf{1}_{\{x_1 \geq h\}} + p_2 \mathbf{1}_{\{x_2 \geq h\}} + p_3 \mathbf{1}_{\{x_3 \geq h\}}$$

under the condition

$$u(x_1, x_2, x_3) \equiv \mathbf{Var} \xi = (p_1 x_1^2 + p_2 x_2^2 + p_3 x_3^2) - (p_1 x_1 + p_2 x_2 + p_3 x_3)^2 \leq \sigma^2.$$

Let us keep p_0, p_1, p_2, p_3 fixed, fixed for a while. Next, one can assume that $0 \leq x_1 \leq x_2 \leq x_3$. Note that in the region $x_3 > x_2$,

$$\frac{\partial u}{\partial x_3} = 2p_3(x_3 - (p_1 x_1 + p_2 x_2 + p_3 x_3)) > 0.$$

That is, $u = \mathbf{Var} \xi$ is an increasing function of x_3 in $[x_2, +\infty)$. But, as a function of x_3 , J is constant in $[0, h)$ and constant $[h, +\infty)$. Hence, any $x_3 \in [0, h)$ can be replaced by $x_3 = x_2$, and any $x_3 \in [h, +\infty)$ can be replaced by $x_3 = \max(x_2, h)$. In the first case, a new random variable ξ will have a smaller variance, the value of J will be unchanged, but ξ will take at most three values: $0, x_1, x_2$. In the second case, the same is true provided that $x_2 \geq h$, and ξ will take the values $0, x_1, x_2, h$ if $x_2 < h$. Now, let $x_2 \geq h$, thus one can put $x_3 = x_2$. Then, by the same type of reasoning, x_2 can be

replaced by $x_2 = \max(x_1, h)$. If $x_1 \geq h$, then ξ will take only the values 0 and x_1 , and once more x_1 can be replaced by h .

The arguments above show that only the case $0 \leq x_1, x_2 < x_3 = h$ need to be worked out and in fact the value $x_3 = h$ may disappear if $p_3 = 0$

Step 4 (reduction to the three atoms case, one of them being h).

Proving (2.6) is thus reduced to study random variables ξ such that

$$\mathcal{L}(\xi) = p_0\delta_0 + p_1\delta_{x_1} + p_2\delta_{x_2} + p_3\delta_h,$$

where $p_0 \geq p$, $p_1, p_2, p_3 \geq 0$, $p_0 + p_1 + p_2 + p_3 = 1$, $0 \leq x_1, x_2 < h$. Now, $J = p_3$ and one needs to maximize the functional J under the condition

$$u(x_1, x_2) \equiv \mathbf{Var} \xi = (p_1x_1^2 + p_2x_2^2 + p_3h^2) - (p_1x_1 + p_2x_2 + p_3h)^2 \leq \sigma^2.$$

Again, p_0, p_1, p_2, p_3 are kept fixed and x_1 and x_2 are allowed to vary in $[0, h)$. In addition, it is assumed that, for p_0, p_1, p_2, p_3 given and fixed, there exists at least one pair $x_1, x_2 \in [0, h)$ such that $u(x_1, x_2) \leq \sigma^2$ (otherwise, the above values of p_0, p_1, p_2, p_3 cannot be considered).

Under the above conditions, $J = p_3$ is constant, and there is nothing to maximize. However, one needs to prove the existence of p_0, p_1, p_2, p_3 such that $u(x_1, x_2) \leq \sigma^2$, for some $x_1, x_2 \in [0, h)$. To finish this step, we only show that if this is at all possible, one may choose $x_1 = x_2$.

First observe that $u = u(x_1, x_2)$ is a positive quadratic form, hence an extremal point of u will also be the global minimum on the whole plane \mathbf{R}^2 . The extremal point satisfies

$$\begin{cases} \frac{1}{2} \frac{\partial u}{\partial x_1} = p_1x_1 - p_1(p_1x_1 + p_2x_2 + p_3h) = 0 \\ \frac{1}{2} \frac{\partial u}{\partial x_2} = p_2x_2 - p_2(p_1x_1 + p_2x_2 + p_3h) = 0 \end{cases}$$

that is, assuming that $p_1, p_2 > 0$,

$$\begin{cases} x_1 = p_1x_1 + p_2x_2 + p_3h \\ x_2 = p_1x_1 + p_2x_2 + p_3h \end{cases}$$

from which it follows that

$$x_1 = x_2 = \frac{p_3}{1 - p_1 - p_2} h. \quad (2.7)$$

Now, since $p_3 < 1 - p_1 - p_2 = p_0 + p_3$, then $x_1 = x_2 \in [0, h)$. Therefore, the global minimum (x_1, x_2) of u on \mathbf{R}^2 is also a global minimum of u on the square $[0, h) \times [0, h)$. Therefore, there exists $(x_1, x_2) \in [0, h) \times [0, h)$ such that $u(x_1, x_2) \leq \sigma^2$, if and only if $u(x_1, x_2) \leq \sigma^2$ with $x_1 = x_2$ defined by (2.7). When, $p_1 = 0$ and/or $p_2 = 0$ we, in fact, deal with ξ taking at most three values.

Step 5: the case $F = p_0\delta_0 + p_1\delta_x + p_2\delta_h$, $0 \leq x < h$, $p_0 \geq p$, $p_1, p_2 \geq 0$, $p_0 + p_1 + p_2 = 1$.

Again, $J = p_2$ is constant. According to (2.7), (changing the notations there), for fixed p_0, p_1, p_2 , the minimal value of $\mathbf{Var} \xi$ as a function of x on $[0, h)$ is attained at

$$x = \frac{p_2}{p_0 + p_2}h.$$

For this value of x , we find

$$\begin{aligned} \mathbf{Var} \xi &= (p_1x^2 + p_2h^2) - (p_1x + p_2h)^2 \\ &= \left(p_1 \frac{p_2^2}{(p_0 + p_2)^2} h^2 + p_2h^2 \right) - \left(p_1 \frac{p_2}{p_0 + p_2} h + p_2h \right)^2 \\ &= \left[\frac{p_2}{(p_0 + p_2)^2} (p_1p_2 + (p_0 + p_2)^2) - \frac{p_2^2}{(p_0 + p_2)^2} (p_1 + (p_0 + p_2))^2 \right] h^2 \\ &= \frac{p_2}{(p_0 + p_2)^2} [p_1p_2 + (p_0 + p_2)^2 - p_2] h^2 \\ &= \frac{p_0p_2}{p_0 + p_2} h^2. \end{aligned}$$

Now, we have to maximize $J = p_2$ under the condition

$$\mathbf{Var} \xi = \frac{p_0p_2}{p_0 + p_2} h^2 \leq \sigma^2. \quad (2.8)$$

From (2.8), when p_0 decreases, $\mathbf{Var} \xi$ also decreases, while $J = p_2 = 1 - p_0 - p_1$ increases (p_1 is fixed). Hence, in the sequel, it is enough to only consider the case $p_0 = p$. The possible maximal value $p_2 = 1 - p$ satisfies (2.8) if and only if $p(1 - p) \leq \sigma^2/h^2$, that is, if and only if $h \leq h(p, \sigma)$. Otherwise, if $h > h(p, \sigma)$, or even if $h = h(p, \sigma)$, the maximal value of $J = p_2$ is, according to (2.8), the value which satisfies

$$\frac{pp_2}{p + p_2} h^2 = \sigma^2.$$

The only solution to this equation is given by

$$p_2 = \frac{p\sigma^2}{ph^2 - \sigma^2}.$$

Lemma 6 follows.

Proof of Corollary 4.

Let $p = \mu(A)$, $q = \mu(B)$. If $p = 0$ or $q = 0$, there is nothing to prove. If $p + q > 1$, then $A \cap B \neq \emptyset$, so $d_1(A, B) = 0$. Thus, we need only consider the case $p + q \leq 1$. Let $p, q > 0$, $p + q \leq 1$, and assume $A \cap B = \emptyset$. Note that

$$h \equiv \sigma \sqrt{\frac{1}{p} + \frac{1}{q}} > h(p, \sigma).$$

Therefore, by (1.2),

$$1 - \mu(A^h) \leq \frac{p\sigma^2}{ph^2 - \sigma^2} = q,$$

and again by (1.2), for all $h_1 > h$, $1 - \mu(A^{h_1}) < q$. Hence, $B \cap (A^{h_1} \setminus A) \neq \emptyset$, and therefore, $d_1(A, B) \leq h_1$. Letting $h_1 \rightarrow h$, completes the proof.

References

- [F] X.Fernique. Régularité de fonctions aléatoires non gaussiennes. *Lecture Notes in Math.* 976 (1981), 2–74.
- [H] L.H.Harper. Optimal numbering and isoperimetric problems on graphs. *J. Comb. Theor.* 1 (1966), 385–393.
- [L] M.Ledoux. Isoperimetry and Gaussian Analysis. *École d'été de Probabilités de Saint-Flour.* (1994).
- [T1] M.Talagrand. An isoperimetric theorem on the cube and the Khinchine–Kahane inequalities. *Proc. Amer. Math. Soc.* 104 (1988), 905–909.
- [T2] M.Talagrand. Concentration of measure and isoperimetric inequalities in product spaces. Preprint (1994).

Serguei. G. Bobkov
Department of Mathematics
Syktyvkar University
167001 Syktyvkar
Russia

Christian Houdré
Center for Applied Probability
School of Mathematics
Georgia Institute of Technology
Atlanta, GA 30322, USA